

Commun. Math. Phys.
81, 239–241 (1981)

Lax Representation for the Systems of S.Kowalevskaya Type

A.M. Perelomov

Institute of Theoretical and Experimental Physics, 117259 Moscow, USSR

Abstract

We describe a number of dynamical systems that are generalizations of the S. Kowalevskaya system and admit the Lax representation.

It is well known (see for example [1,2]) that the equations of motion of a three-dimensional heavy rigid body rotated about a fixed point is a completely integrable dynamical system only in the cases of Euler [3], Lagrange [4] and Kowalevskaya [5]. In the present note we describe a number of systems that are of the Kowalevskaya type and admit the Lax representation¹.

1. Let $\mathcal{G}$ be the Lie algebra of a group G , $\mathcal{G}^*$ be the space dual to $\mathcal{G}$ and $\{x_\alpha\}$ be the coordinates of a point in the space $\mathcal{G}^*$. In the space $\mathcal{F}(\mathcal{G}^*)$ of smooth functions on $\mathcal{G}^*$, let us define the Poisson bracket ²

$$\{f, g\} = C_{\alpha\beta}^\gamma x_\gamma \partial^\alpha f \partial^\beta g, \quad \partial^\alpha = \partial / \partial x_\alpha. \quad (1)$$

Here $C_{\alpha\beta}^\gamma$ are the structure constants of the Lie algebra. The space $\mathcal{F}(\mathcal{G}^*)$ is endowed by the above formula with the structure of the Lie algebra. A

¹ Note that the Lax representation for equations of motion of a rigid body was discovered first by Manakov [8] for some particular cases.

² In this note we use the tensor notations; in particular, everywhere repeated indices imply summation.

dynamical system in $\mathcal{G}^*$ is determined by a Hamiltonian function $H(x) \in \mathcal{F}$, so that the equations of motion have the form

$$\dot{x}_\alpha = \{H, x_\alpha\}. \quad (2)$$

The coadjoint representation of the group G acts on the space $\mathcal{G}^*$. Orbits of this representation are invariant with respect to an arbitrary Hamiltonian H , and are the phase spaces of the considered systems.

2. Let G be a compact simple Lie group, K be its subgroup such that the factor space G/K is a symmetric space [6]. Then $\mathcal{G} = \mathcal{K} \oplus \mathcal{S}$, $\mathcal{K}$ is the Lie algebra of the group K , $\mathcal{S}$ is the orthogonal complement to $\mathcal{K}$ in $\mathcal{G}$ relative to the Killing-Cartan form.

It is well known that a certain irreducible representation $T^0(k)$ of the group K acts on the space $\mathcal{S}$. Let us consider an irreducible representation $T(g)$ of the group G , such that under the restriction of this representation on the subgroup K , the irreducible representation $T^0(k)$ is contained in $T(g)$ with the unit multiplicity. Let V be a vector space in which the representation $T(g)$ acts. Then it is possible to define the group $\tilde{G} = G \cdot V$ that is a semidirect product of the group G and the abelian vector group V . Let us denote by V^0 the subspace of V on which the representation $T^0(k)$ acts, and by A the projection operator on the subspace V^0 . Let V^1 be the orthogonal complement to V^0 : $V = V^0 \oplus V^1$. Note that the spaces V^0 and $\mathcal{S}$ are isomorphic because $T(g)$ contains $T^0(k)$ only once. Therefore we have the decomposition

$$\tilde{\mathcal{G}} = \mathcal{K} \oplus \mathcal{S} \oplus \mathcal{V}^0 \oplus \mathcal{V}^1,$$

and the analogous decomposition for the space $\tilde{\mathcal{G}}^*$:

$$\tilde{\mathcal{G}}^* = \mathcal{L} \oplus \mathcal{N} \oplus \mathcal{P} \oplus \mathcal{T}. \quad (3)$$

Here $\mathcal{L} = \mathcal{K}^*$, $\mathcal{N} = \mathcal{S}^*$, $\mathcal{P} = \mathcal{V}^{(0)*}$, $\mathcal{T} = \mathcal{V}^{(1)*}$. In addition, $\dim \mathcal{N} = \dim \mathcal{P} = n$, and $\mathcal{N}$ and $\mathcal{P}$ are isomorphic relative to the action of the group K .

Let $\hat{l}$ be the matrix of the representation T of the Lie algebra $\mathcal{G}$, $\hat{l} = \{l_{jk}\}$ corresponds to the matrix of the representation T^0 of the algebra $\mathcal{K}$ acting in the space $\mathcal{P}$. Let us introduce the matrix, which is important in the sequel,

$$L = A \left[-\hat{l}^2 + (\gamma \otimes p + p \otimes \gamma) \right] A, \quad (4)$$

where $p \in \mathcal{P} \oplus \mathcal{T}$ and $\gamma \in \mathcal{P}$ is the constant vector in $\mathcal{P}$. We identify $\mathcal{P}$ and $\mathcal{P}^*$ by means of the K -invariant scalar product on $\mathcal{P}$.

The system of S. Kowalevskaya type is defined by the Hamiltonian

$$H = \frac{1}{2} \operatorname{tr} L = \alpha I_2(l) + \beta \mathcal{J}_2(n) + (\gamma p). \quad (5)$$

Here $I_2(l)$ and $\mathcal{J}_2(n)$ are the quadratic functions on $\mathcal{L}$ and $\mathcal{N}$ respectively that are invariant relative to the coadjoint representation of the group K and $p \in \mathcal{P}$.

3. Let us consider in detail the case: $G = SO(n+1)$, $K = SO(n)$, $\tilde{G} = G \cdot V = E(n+1)$ is the motion group of the $(n+1)$ -dimensional Euclidean space $V = \mathbb{R}^{n+1}$, $V^0 = \mathbb{R}^n$, $V^1 = \mathbb{R}^1$. Let $\hat{l}_{jk} = -\hat{l}_{kj}$ and $\hat{p}_m$ ($j, k = 1, \dots, (n+1)$) be the standard basis in the space of linear functions on $\mathcal{G}^*$ and $\mathcal{V}^*$ respectively with the standard Poisson brackets.

The Hamiltonian (5) takes now the form

$$2H = 2 \sum_{j < k}^n l_{jk}^2 + \sum_{j=1}^n n_j^2 + 2 \sum_{j=1}^n \gamma_j p_j. \quad (6)$$

Theorem. *The equations of motion (2) of the system with Hamiltonian (6) are equivalent to the Lax equation*

$$\dot{L} = [L, M], \quad (7)$$

where L is given by the formula (4), and $M = cl = c\hat{A}lA$, c is a constant.

The theorem is verified by a direct calculation.

From the Lax representation (7), it follows immediately that the eigenvalues of the matrix L or the quantities $I_{2k} = k^{-1} \operatorname{tr}(L^k)$, $k = 1, \dots, n$ are integrals of motion for Eqs. (2). Notice $I_2 = 2H$.

There are also $(n-1)(n-2)/2$ linear integrals related to the invariance of H with respect to the group $G_0 = SO(n-1)$ that leaves the vector γ invariant, and also $([n/2] + 1)$ independent polynomials that are invariant with respect to the coadjoint representation of the group $\tilde{G}$. Using the theory of reduction of hamiltonian systems with symmetries [2,7] we may show that the above system is reduced to a system with $2n$ -dimensional phase space. It can be verified that integrals $I_2, I_4, \dots, I_{2n}$ are functionally independent and in involution. Therefore, the systems under consideration are completely integrable.

Finally, we wish to note the following facts:

1. The system considered by S. Kowalevskaya corresponds to $n = 2$.
2. Similar results are valid for other symmetric spaces. The simplest ones are the cases of the spaces of rank one: $SU(n + 1)/SU(n) \times U(1)$, $Sp(n + 1)/Sp(n) \times Sp(1)$ and $F_4/SO(9)$.
3. If $p \in \mathcal{T}$ in formula (5), then we deal with the generalization of the Lagrange case. In this case, the constants α and β may be arbitrary.
4. The results of this section are valid also for similar systems, related to group $\tilde{G} = SO(n + 2)$.

The detailed presentation of results will be published elsewhere.

Acknowledgement. I am grateful to V.Golo for the improvement of the language of this note.

References

- [1] Motion of a rigid body around a fixed point. A collection of papers, dedicated to the memory of S. Kowalevskaya. Acad. of Sci. USSR, Moscow (1940) (in Russian)
- [2] Arnold, V.I.: Mathematical methods of classical mechanics. New York: Springer 1978
- [3] Euler, L.: Decouverte d'un nouveau principe de Mecanique. Mem. de l'Acad. des Sciences de Berlin 1758
- [4] Lagrange, J.L.: Mecanique analytique . Paris 1788
- [5] Kowalevskaya, S.: Acta Math. **12**, H.2 177–232 (1889)
- [6] Helgason, S.: Differential geometry and symmetric spaces. New York: Acad. Press 1962
- [7] Marsden, J., Weinstein, A.: Rep. Math. Phys. **5**, 121–130 (1974)
- [8] Manakov, S.: Funkt. Anal. Priloz. **10**, 93–94 (1976)